

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 39/29		A1	(11) International Publication Number: WO 95/05194
			(43) International Publication Date: 23 February 1995 (23.02.95)
(21) International Application Number: PCT/GB94/01646		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD).	
(22) International Filing Date: 29 July 1994 (29.07.94)			
(30) Priority Data: 9316745.0 12 August 1993 (12.08.93) GB			
(71) Applicant (for all designated States except US): MEDEVA HOLDINGS B.V. [NL/NL]; Churchill-Laan 223, NL-1078 ED Amsterdam (NL).		Published With international search report.	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): CHATFIELD, Steven, Neville [GB/GB]; Medeva Vaccine Research Unit, Imperial College of Science, Technology and Medicine, Exhibition Road, London SW7 6AY (GB). ROBERTS, Mark [GB/GB]; Medeva Vaccine Research Unit, Imperial College of Science, Technology and Medicine, Exhibition Road, London SW7 6AY (GB).			
(74) Agents: HUTCHINS, Michael, Richard et al.; Fry Heath & Spence, The Old College, 53 High Street, Horley, Surrey RH6 7BN (GB).			

(54) Title: VACCINE COMPOSITIONS



(57) Abstract

The invention provides the use of Hepatitis A virus capsid, or a mucosally immunogenic fragment or epitope thereof, for the manufacture of a mucosal vaccine composition for administration to a mucosal surface of a patient to induce production of serum Immunoglobulin G antibody against Hepatitis A. Preferably the vaccine composition is administered by the intranasal route.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

VACCINE COMPOSITIONS

The present invention relates to vaccine compositions for delivery to mucosal surfaces, and to a method of inducing, in a mammal, an immune response to an antigen by delivering the antigen to a mucosal surface of the mammal. More particularly, the present invention relates to vaccine compositions for inoculating a mammal such as a human against picornavirus infection and particularly Hepatitis A infection.

Hepatitis A is an acute disease caused by infection with a small picornavirus closely related to the poliovirus. Infection is spread by the faecal/oral route and consequently the disease is endemic in areas where hygiene and sanitation standards are low. The risk of travellers to developing countries acquiring Hepatitis A is far greater than that of contracting typhoid and cholera (40 and 800 times respectively).

The virus itself is not directly cytopathic. The liver damage resulting from Hepatitis A virus (HAV) infection arises from destruction of virally infected cells by the host's cytotoxic T-lymphocytes. There is only a single serotype of HAV and infection results in long-term

immunity, characteristics that are ideal for developing Hepatitis A prophylaxis. Protection is mediated by neutralising antibodies that prevent entry of hepatitis A virus into hepatocytes. Passive immunisation with purified human serum γ -globulin provides short term protection against the disease and until recently this was the only means of preventing hepatitis A.

In recent years, HAV vaccines have been developed but development has focused on inactivated and live attenuated vaccines. Both types of vaccines are prepared from HAV propagated in tissue culture cells. HAV replication is slow and the majority of the virus remains cell associated, and consequently the viral yields are low and relatively commercially unattractive. The problem of low viral yield could be overcome by using recombinant techniques which allow for the production of large quantities of proteins. However, it is important to ensure correct processing and folding of HAV proteins because the known neutralising epitopes are conformationally dependent. It has proved difficult so far to obtain recombinant HAV antigens that elicit appropriate immune responses.

One recombinant HAV antigen that has proved successful in inducing protection against HAV when injected parenterally is the HAV capsid preparation developed by American Biogenetic Sciences. American Biogenetic Sciences have succeeded in producing empty HAV capsids in eucaryotic cells using vaccinia and baculovirus expression vectors. The recombinant capsids are recognised by neutralising

monoclonal antibodies, induce protection against HAV in chimpanzees when injected parenterally and are produced in considerably larger quantities than that obtained by conventional means. The HAV capsids are disclosed in International Patent Application WO-A-9301279, the disclosure in which is incorporated herein by reference.

A disadvantage with many vaccination regimens is that it is frequently necessary to administer the vaccine composition by means of injection, a factor which has a potential deterrent effect to many people, particularly when follow-up or booster injections are required to complete a course of treatment. One way of overcoming this problem would be to administer the vaccine composition to the oral or nasal mucosa, but although immunisation by the oral or intranasal route has been explored with certain other antigens, it has been found usually to be less effective in evoking serum antibodies than parenteral immunisation. For example, the article by M.H. Sjogren *et al*, Vaccine, Vol. 10, Suppl. 1, S135-S137, 1992, describes the administration of a live attenuated hepatitis A vaccine by either the oral route or the intramuscular route. Whereas intramuscular administration elicited a good serum antibody response, an antibody response to oral administration was not observed at any dose.

The fact that there are very few mucosal vaccines commercially available indicates that there are problems with developing such vaccines. May non-living soluble antigens, particularly those used traditionally by

immunologists, such as ovalbumin (OVA) and Keyhole Limpet Haemocyanin (KLH) are poor mucosal immunogens. Large doses of such antigens are necessary to induce any responses but large doses can also cause tolerance in the individual to subsequent parenteral exposure to antigen, a condition known as Oral Tolerance. Although some microbial components such as the cholera toxin (CT) or E.coli heat labile toxin (LT) or the non-toxic binding portions of these toxins (CT-B and LT-B) have been found to be potent mucosal immunogens eliciting strong secretory and circulating antibodies, the reasons why such molecules are good mucosal immunogens has not been fully elucidated. One property that may be important is the ability of these molecules to bind to mucosal epithelial cells via certain surface receptors, although it has been found in studies by others that there is not necessarily a correlation between the ability of an antigen to bind to eucaryotic cells and its mucosal immunogenicity. In the present case, it is not known, at the molecular or cellular level, how HAV enters the body, nor is it known whether specific receptors are involved.

Thus, as far as we are aware, there is currently no way of predicting with any certainty whether a given antigen will possess good mucosal immunogenicity.

It has now been found that the recombinant empty HAV capsid referred to hereinabove, when administered mucosally, and in particular intranasally, is efficient at inducing serum anti-HAV antibodies. Thus, when the HAV

capsid preparation was administered intranasally, following first and second booster doses, seroconversion to anti-HAV was observed in the majority of animals, and this compared favourably with the administration of the antigen in the presence of an alum adjuvant by the subcutaneous route.

Accordingly, in a first aspect, the invention provides the use of Hepatitis A virus capsid, or mucosally immunogenic fragments or epitopes thereof, for the manufacture of a mucosal vaccine composition for administration to a mucosal surface of a patient to induce the production of serum Immunoglobulin G antibody against Hepatitis A virus.

In a second aspect, the invention provides a vaccine composition for application to a mucosal surface, the composition comprising Hepatitis A virus capsid, or a mucosally immunogenic fragment or epitope thereof, and a pharmaceutically acceptable carrier.

In a still further aspect of the invention, there is provided a method of inducing the production of serum Immunoglobulin G antibody against Hepatitis A virus in a host such as a mammal (eg. human), which method comprises administering an effective amount of a Hepatitis A virus capsid antigen, or a mucosally immunogenic fragment or epitope thereof, directly to a mucosal surface in the host.

The mucosal delivery compositions of the present invention can be formulated, for example, for delivery to one or more of the oral, gastro intestinal, and respiratory (eg. nasal and bronchial) mucosa.

Where the composition is intended for delivery to the respiratory (eg. nasal or bronchial) mucosa, typically it is formulated as an aqueous solution for administration as an aerosol or nasal drops, or as a dry powder eg. for inhalation.

Compositions for administration as nasal drops may contain one or more excipients of the type usually included in such compositions, for example, preservatives, viscosity adjusting agents, tonicity adjusting agents, buffering agents and the like. The vaccine compositions of the present invention may also take the form of compositions intended to deliver the antigen to mucosal surfaces in the gastro intestinal tract. Such compositions can be provided with means for preventing degradation of the antigens by the gastric juices, for example by encasing the vaccine preparation in a capsule within a protective matrix or coating of known type.

The quantity of Hepatitis virus A capsid administered to the patient typically is selected such that it is non-toxic to the patient at concentrations employed to elicit an immune response. For example, the concentration of capsid administered may lie in the range 0.1mg to 100mg per kg/host.

The invention will now be illustrated in more detail by reference to the specific embodiments described in the following examples, and illustrated in the accompanying drawings. The examples are intended to be purely illustrative of the invention and are not intended to limit

its scope in any way.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates the individual and mean serum anti-HAV titres following three doses of HAV capsids by each of the intranasal, oral and subcutaneous routes.

A sample of recombinant HAV capsids was obtained from American Biogenetic Sciences, Reyniers Germ Free Building, P.O. Box 1001 Notre Dame, Indiana 46556, USA. The sample was prepared from vaccinia-HAV infected Vero cells. Cells were lysed with NP40 and then extracted with trichlorotrifluoroethane. The aqueous phase was concentrated and then chromatofocussed on Biogel A-15 column. Fractions containing empty capsids were pooled and concentrated. Formalin was added to inactivate any remaining vaccinia virus. The content of HAV capsids in the sample as given below is expressed in terms of ELISA units (EU). The EU values have been standardised on a sample of Hepatitis A virus obtained from SmithKline Beecham. The sample received contained 52 ELISA units (EU) HAV capsids per μ l. The protein content of the sample was 30 mg/ml of which 100ng/ml was estimated to be HAV antigen.

Recombinant HAV capsids of the aforementioned type can be prepared in accordance with the methods as set out in International Patent Application WO-A-9301279 (PCT/US92/05714).

Mice were immunised orally and intranasally (I/N) with different doses of HAV. A small quantity (1 µgm) of cholera toxin (CT) was included in the material given to some groups of mice to act as an adjuvant. CT was used because it is the most potent mucosal adjuvant known. A separate group of mice were immunised parenterally with HAV adsorbed to aluminium hydroxide as an adjuvant as a positive control. The groups were as set out in Table 1 below:

TABLE 1

Group	Number of Mice	Dose Elisa Units (EU)	Adjuvants
1. Parenteral	5	250 x 3	Alum
2. Parenteral Control	5	-	Alum
3. Oral	10	500 x 3	-
4. Oral	10	250 x 3	-
5. Oral	10	125 x 3	-
6. Oral	10	250 x 3	CT ^a
7. Oral Control	5	-	CT
8. I/N ^b	10	-	-
9. I/N	10	500 x 3	-
10. I/N	10	250 x 3	-
11. I/N	10	125 x 3	CT
12. I/N Control	5	250 x 3	CT
		-	
^a CT, Cholerae toxin (1µg/dose) ^b I/N, intranasal			

DayProcedure

0
20

Primary immunisation
Sample bleed

24	1st booster immunisation
31	Bleed, gut and nasal washes
47	2nd booster immunisation
54	Sample bleed

The immunisation was carried out by the method set out in Table 2 below:

TABLE 2

IMMUNISATION DETAILS

Route	Volume Delivered (µl)	Diluent	Apparatus	Anaesthetic (Halothane)
Intranasal	30	PBS	micro pipette	light
Oral	200	5% Bicarbonate sol.	gavage needle	light
Subcutaneous	100	PBS	needle	nil

Anti-HAV responses were analysed using a capture ELISA technique as follows. Human convalescent polyclonal serum from an individual with known Hepatitis A was coated on 96 well plastic plates. The polyclonal serum captures HAV capsids binding them to the plate. Mouse serum was then incubated with the HAV capsid bound plates and the mouse serum reactivity to HAV determined using labelled anti-mouse antibodies. The protocol for the assay was as follows:

PROTOCOL (All volumes 50 µL/well unless otherwise stated)

- 1) Coat Costar EIA plates (Cat no: 3590) overnight, 4°C, with 1:25000 human capture antibody diluted in PBS.
- 2) Wash plate x3 with phosphate buffered saline/Tween (0.05%) (PBST).

10

- 3) Block 1hr, 37°C, 1%BSA(Sigma, Cat no: A7888) in PBST(200µl/well).
- 4) Wash plate x3 PBST.
- 5) Coat plates with sample containing HAV (0.1EU/µl) in PBST (0.1%BSA), 2-3 hours 37°C.
- 6) Wash plate x3 PBST.
- 7) Incubate plate with mouse serum diluted PBST, 2-3 hours, 37°C.
- 8) Wash plate x3 PBST.
- 9) Incubate, 1-2 hours, 37°C, with anti-mouse IgG, 1:1000 (goat, Sigma, Cat No: B7022) in PBST.
- 10) Wash plate x3 PBST.
- 11) Incubate, 1-2 hours, 37°C, with Streptavidin-peroxidase 1:1000 (Dako, Cat No: P397) in PBST.
- 12) Wash plate x3 PBST.
- 13) Add substrate, OPD in phosphate-citrate buffer (Sigma, Cat no: P8287), incubate for up to 30 mins 37°C.
- 14) Read colour development after stopping substrate reaction (3M H_2SO_4).

Human sera and HAV capsid sample supplied by ABS.

Optimum conditions of capture Antibody and HAV capsid sample were selected to minimise the S/N ratio.

Problems of high background noise, attributed to reactivity of the mouse serum against the human sera capture antibody were encountered. Reduction of the background was obtained by diluting out the capture antibody, beyond the dilution recommended by ABS.

Similar results for capsid capture were obtained using, 1:1000-1:25000 dilutions of the capture Antibody.

The serum IgG anti-HAV responses were as shown in Table 3 below:

TABLE 3

Table 3. Mean serum IgG anti-HAV response following a primary and booster immunisations

Dose	Intranasal				Titre				Sub/cut
	125	250	500	250+CT	125	250	500	250+CT	250+Alum
1	<50	<50	50	6000	<50	<50	<50	<50	6000
2	<250	250	9462	8772	<250	<250	<250	<250	10438
3	7650	3777	23030	46250	<250	250	480	2170	54000

Mean titre calculated from responding mice only

As can be seen from the Table, after a single dose, anti-HAV antibodies could be detected in the sera of mice immunised subcutaneously with 250EU and intranasally with 500EU. There was no detectable response in any of the other groups. Boosting greatly enhanced the response in the subcutaneous and intranasal 500 groups, giving titres of approximately 10500 and 9500 respectively. Also, a very similar response was seen in mice receiving 250EU plus CT I/N and a low but measurable response was detected in the sera of half the mice receiving two doses of 250EU I/N. No serum response was detectable in any of the orally immunised mice at this point. Further boosting did result in seroconversion of some of the orally immunised mice into 250EU, 500EU and 250EU + CT groups. The response was

greatest in the later group. After three doses, all of the I/N immunised mice had seroconverted, the magnitude of the response being dose dependant in the absence of CT. I/N immunisation with 500EU, 250EU+CT produced high serum titres that were slightly lower but comparable to those produced by S/C immunisation with 250EU, HAV adsorbed to alum.

Figure 1 illustrates the individual and mean serum anti-HAV titres following three doses of HAV capsids. As can be seen, CT greatly augmented the serum anti-HAV response of I/N and orally administered antigen. The titres in mice receiving 250EU+CT I/N were more than 10-fold rated at each time point than those evoked by I/N immunisation with 250EU alone. There was a similar difference in anti-HAV titre in the mice immunised orally with 250EU with and without CT. CT also increased the number of mice seroconverting. Moreover, the titres in the I/N 250EU+CT group follow each immunisation were very similar to those of mice given 250EU adsorbed to alum and given subcutaneously.

Table 2 below shows the number of mice in the different groups seroconverting following the first and second booster doses. Seroconversion was dose and route dependant. Subcutaneous immunisation with 2 doses of 250EU led to seroconversion in all the mice, whereas the same dose, only half of the mice to whom the antigen had been administered intranasally, and none of the mice to whom antigen had been administered orally, had seroconverted.

After the second boost, all of the mice in the I/N groups had seroconverted, including those in the 125EU dose group that exhibited no response after two doses. Likewise, the orally immunised mice started a response after the second boost although even the addition of CT did not result in the seroconversion of all of the mice.

TABLE 4

Rate of seroconversion to anti-HAV following the first and second booster immunisations

Dose	Intranasal			Oral			Sub/cut		
	125	250	500	250+ CT	125	250	500 CT	250+	250+Alum
2	0/4*	2/4	4/4	4/4	0/4	0/4	0/4	0/4	5/5
3	6/6	5/5	5/5	5/5	0/4	1/4	2/4	3/4	5/5

* No. of mice responding/No. of mice tested

Table 4 shows the number of mice in the different groups seroconverting following the first and second booster doses. Seroconversion was dose and route dependant. subcutaneous immunisations with two doses of 250EU led to seroconversion in all the mice, whereas at the same dose only half of the mice treated intranasally and none of the mice treated orally had seroconverted. After the second boost all of the mice in the intranasal groups had seroconverted, including those in the 125EU dose group that exhibited no response after 2 doses. Likewise the orally immunised mice started to respond after the second boost

although even the addition of CT did not result in the seroconversion of all of the mice.

The aforementioned examples are given by way of illustration only and are not intended the scope of the application, which is limited only by the claims appended hereto.


CLAIMS

1. The use of Hepatitis A virus capsid, or a mucosally immunogenic fragment or epitope thereof, for the manufacture of a mucosal vaccine composition for administration to a mucosal surface of a patient to induce production of serum Immunoglobulin G antibody against Hepatitis A.
2. The use according to Claim 1 wherein the Hepatitis A virus capsid is recombinant.
3. A method of inducing the production of serum Immunoglobulin G antibody against Hepatitis A virus in a host such as a mammal (eg. human), which method comprises administering an effective amount of Hepatitis A virus capsid, or a mucosally immunogenic fragment or epitope thereof directly to a mucosal surface in the host.
4. A method according to Claim 3 wherein the Hepatitis A virus capsid is recombinant.
5. The use according to Claim 1 or Claim 2 wherein the mucosal vaccine composition is packaged for delivery to the nasal or bronchial mucosa.
6. The use according to Claim 4 wherein the composition

is formulated as an aqueous solution for administration as an aerosol or nasal drops, or as a dry powder for inhalation.

7. The use according to Claim 1 or Claim 2 wherein the composition is adapted for oral or gastro-intestinal delivery of the Hepatitis A virus capsid.

1/1



INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 94/01646

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K39/29

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO,A,94 17827 (LYFJATHROUN H.F. ET AL.) 18 August 1994 see page 8, line 1 - page 10, line 17 ---	1,3,5,6
X	WO,A,93 01279 (AMERICAN BIOGENETIC SCIENCE INC.) 21 January 1993 cited in the application see page 18, line 5 - page 23, line 22 ---	1-7
X	DATABASE WPI Section Ch, Week 9129, Derwent Publications Ltd., London, GB; Class B04, AN 91-212580 & JP,A,3 135 923 (KAGAKU OYOBI KESSEI) 10 June 1991 see abstract --- -/--	1,3,5,6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

11 November 1994

Date of mailing of the international search report

24. 11. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Rempp, G

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/GB 94/01646

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>EP,A,0 418 626 (TAKEDA CHEMICAL INDUSTRIES, LTD.) 27 March 1991 see column 2, line 38 - column 3, line 12 see column 7, line 21 - line 52 -----</p>	1-7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 94/01646

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9417827	18-08-94	NONE	
WO-A-9301279	21-01-93	US-A- 5294548	15-03-94
		AU-A- 2371692	11-02-93
EP-A-0418626	27-03-91	DE-D- 69005572	10-02-94
		DE-T- 69005572	07-04-94
		JP-A- 3178995	02-08-91